



ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

Current Opinion in
Neurobiology

Nanoscale analysis of structural synaptic plasticity

Jennifer N Bourne and Kristen M Harris

Abstract

Structural plasticity of dendritic spines and synapses is an essential mechanism to sustain long lasting changes in the brain with learning and experience. The use of electron microscopy over the last several decades has advanced our understanding of the magnitude and extent of structural plasticity at a nanoscale resolution. In particular, serial section electron microscopy (ssEM) provides accurate measurements of plasticity-related changes in synaptic size and density and distribution of key cellular resources such as polyribosomes, smooth endoplasmic reticulum, and synaptic vesicles. Careful attention to experimental and analytical approaches ensures correct interpretation of ultrastructural data and has begun to reveal the degree to which synapses undergo structural remodeling in response to physiological plasticity.

Address

Center for Learning and Memory, Department of Neurobiology,
University of Texas, Austin, TX 78712-0805, USA

Corresponding author: Harris, Kristen M (kharris@mail.clm.utexas.edu)

Current Opinion in Neurobiology 2011, 22:1–11

This review comes from a themed issue on
Synaptic structure and function
Edited by Morgan Sheng and Antoine Triller

0959-4388/\$ – see front matter

© 2011 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.conb.2011.10.019

In the 1950s, transmission electron microscopy was first used to reveal the diversity in synaptic structure and composition in the central nervous system [1,2]. Since then, visualization and reconstruction of serial thin sections have provided three-dimensional contexts in which to understand how synapses are modified with plasticity, learning, and sensory input [3,4^{••},5–16,17^{••}]. Three-dimensional reconstruction from serial section electron microscopy (ssEM) has proven invaluable for the comprehensive analysis of structural synaptic plasticity. It has provided the needed nanometer resolution to localize and measure key subcellular structures, such as the postsynaptic density (PSD) and presynaptic vesicles which define a synapse, polyribosomes as sites of local protein synthesis, smooth endoplasmic reticulum (SER) for local regulation of calcium and trafficking of membrane proteins, endosomes for recycling, and fine astroglial processes at the perimeter of some synapses. Thus, ssEM

is an essential tool for nanoscale analysis of the cell biological and anatomical modifications that underlie changes in synaptic strength. Here we discuss several important issues associated with interpreting the functional significance of structural synaptic plasticity, especially during long-term potentiation, a widely studied cellular model of learning and memory.

General principles

Successful interpretation of structural synaptic plasticity using electron microscopy requires consideration of multiple factors. Especially important are the methods of tissue preservation, experimental preparation, LTP induction, and appropriate analysis.


Tissue preservation

Rapid freezing, particularly combined with high-pressure, followed by freeze-substitution is considered to be the best method for preserving extracellular space, membrane structure, and molecular composition of the cytoplasm [18[•],19]. However, the amount of tissue with high quality freezing is limited to small patches located unpredictably throughout the region of interest. Outside these tiny patches of high quality preservation, the tissue is grossly distorted by ice-crystal artifact. Hence, this approach has not been widely adopted for experimental studies of ultrastructural synaptic plasticity where complete access to whole regions of interest is needed to trace dendrites and axons for sufficient length and volume to investigate changes in circuitry and structure. Currently, the gold standard for tissue preservation is rapid *in vivo* perfusion of fixatives at specific times following the induction of plasticity or consolidation of behavioral tasks (Figure 1a) [12–14,16,17^{••}]. However, functional synaptic plasticity is widely studied in acute or organotypic brain slices and tissue culture where electrophysiological, genetic, or pharmacological manipulations can be restricted to specific regions of interest. In these preparations, perfusion-quality tissue preservation is routinely obtained by microwave-enhanced rapid fixation followed by routine tissue processing (Figure 1b) [3,4^{••},5,6,10]. Recent findings also suggest that microwave-enhanced fixation and processing by freeze-substitution produce high quality images [18[•]].

Experimental preparation of brain slices

Although the brain slice preparation has proven invaluable for physiological studies of long-term potentiation (LTP) [20], one must be aware that differences in slice preparation and the recovery time before experimentation can have profound effects on baseline ultrastructure. For example, upon dissection of the hippocampus

Figure 2



Temperature during slice preparation ultimately influences total synapse number in the recovered slices. Tissue quality of hippocampal neuropil in stratum radiatum of area CA1 is comparable across (a) slices prepared under ice-cold conditions and maintained at $\sim 32^{\circ}\text{C}$ for 9–10 hours in an interface chamber, (b) hippocampus perfusion-fixed *in vivo*, and (c) in slices prepared at room temperature and maintained at $\sim 32^{\circ}\text{C}$ for 4.5–5.5 hours in an interface chamber. (d) Three-dimensional reconstructions of dendrites from each of the three conditions. Scale cube = 1 μm on a side. (e) Dendrites from ice-cold slice preparations had a higher spine density at 9–10 hours *in vitro*, compared to perfusion-fixed hippocampus and room temperature slices (** $P < 0.02$).

Adapted from Bourne *et al.* [31].

and provide a better baseline from which to detect plasticity based changes in ultrastructure.

LTP induction paradigm


Another important consideration for evaluating LTP-mediated ultrastructural plasticity of synapses is the induction paradigm. Early studies used tetanic stimulation *in vivo* (100 Hz for 1 s) followed by instant brain removal and rapid freezing of a chopped surface and found evidence for spine swelling within minutes after the stimulation compared to control brains [32,33]. Others used tetanic stimulation and perfusion of chemical fixatives *in vivo* [34–36] or immersion fixation of brain slices [37], with limited success, although the induction paradigm was confounded by the limitations of single-section analyses discussed below. Brain slices have also been exposed to a cocktail of substances, to produce chemical LTP, which presumably enhances synaptic strength at all

synapses in the slice, but surprisingly little net change in synapse ultrastructure was detected relative to control slices [8].

Theta burst stimulation (TBS) engages multiple plasticity mechanisms including back propagating action potentials, release of neurotrophic factors, activation of voltage gated calcium channels, and enhancement of presynaptic release; all of which are absent during tetanic stimulation [38–40]. TBS also better resembles the endogenous neuronal firing patterns in the hippocampus [41]. Induction of LTP with TBS in area CA1 of mature animals resulted in a pronounced turnover of synapses, protein synthesis-mediated enlargement of spines and synapses, and a coordinated loss of small spines that was perfectly counterbalanced by an enlargement of the remaining synapses [4**]. As discussed below, these findings suggest that closer mimicking of the *in vivo* firing

4 Synaptic structure and function

Figure 3



Serial images through two spines illustrate the need for ssEM to identify the location of PSDs and polyribosomes. (a–c) Serial images illustrating the presence (black arrows) of polyribosomes on some sections and their absence in the same locations on adjacent sections (white arrows). (ai–ci) Same images with dendritic spines highlighted in yellow, PSDs in red and polyribosomes in magenta. Scale bar = 0.5 μm for both columns. (d) Three-dimensional reconstruction of spines (yellow), PSDs (red) and polyribosomes (magenta). Scale cube = 0.5 μm on a side.

patterns results in more robust structural synaptic plasticity not previously found with tetanic induced or chemically induced LTP in brain slices.

Analysis of ultrastructural synaptic plasticity

Sampling methods in EM can markedly influence the detection of structural synaptic plasticity. Complete reconstruction of all objects in the local neuropil provides the most precise quantification of changes in connectivity (number), size and composition of spines, synapses and associated structures. This approach, however, is very time-consuming [42]. Computer-assisted image analysis to automate alignment and segmentation of objects through ssEM sections are being developed and hold promise for future application [43^{••},44^{••},45]. In the past, single-section analyses have dominated the field with limited results for two reasons: first because it is often impossible to identify a synapse or other structures on a single EM section (Figure 3), and second, because counts are biased due to differences in sizes of synapses and other structures, such that larger objects are more likely to be counted but less likely to be accurately identified. For example, single-section analyses have suggested more shaft and sessile spines form after induction of LTP in hippocampal area CA1 [36,37], although it is impossible to distinguish reliably the spiny dendrites of CA1 pyramidal cells from the non-spiny dendrites of inhibitory interneurons on individual sections used for those analyses. Stereological studies using a small number of serial EM sections have suggested an increase in the frequency of perforated synapses [46] and an increase in spine dimensions following the induction of LTP in the mature dentate gyrus [32,33,47]. Similarly, single section analyses following hippocampus-dependent learning found only transient [48,49] or subtle [50] changes in the number and morphology of synapses. However, the reported lack of profound changes with learning and plasticity may be confounded by missed and incorrectly identified synapses or a misrepresentation of their identity, dimensions and composition.

Until dense reconstruction of all structures in a potentiated network becomes routine [43^{••},44^{••},45], computer-assisted manual reconstruction of representative dendrites, axons, and perisynaptic astroglial processes in well-defined regions of interest provides the best alternative to evaluate ultrastructural synaptic plasticity. Sparse sampling from a well-defined subpopulation of dendrites ensures that comparable populations are evaluated across experimental conditions.

Sample results from ssEM that illustrate ultrastructural plasticity of synapses

Synapses are dynamic structures surrounded by a complex neuropil including dendrites, axons, and astroglial processes. Synapses within a small area of neuropil can vary greatly in their size, shape, composition of

subcellular organelles and access to perisynaptic astroglial processes. All of these factors influence functional synaptic plasticity, hence it is important to identify their structural relationships accurately. Here we discuss representative examples where ssEM has served to disambiguate processes of synapse formation and growth during synaptic plasticity.

At present there are essentially two strategies amenable to investigating structural synaptic plasticity at the ultrastructural level, each with their strengths and caveats. One approach has been to image dendrites while inducing plasticity in organotypic slice cultures [51] or while eliciting stereotypical behavior in a well-known circuit, such as whisking and mouse barrel cortex [12–14]. Coincident changes in dendritic spine structure and physiology or behavior are tracked and approximate regions of interest are noted by gross landmarks such as blood vessels. The tissue is fixed and previously visualized cells are processed to visualize an electron dense marker, such as HRP enhanced antibodies to GFP. ssEM is then used to identify synapse locations on the previously visualized cells. An important caveat is that the electron dense product often obscures subcellular components that help to understand the local cell biology of the synapses under investigation.

The other approach has been to stimulate most or all synapses with electrical or pharmacological manipulations and perform statistics by comparing populations of synapses that underwent the differential activation [4^{••},6–8,10,52–54]. This approach allows for a larger sample size but has the caveat of not knowing which individual synapses were potentiated. In addition, pre-synaptic and postsynaptic organelles, molecules, and electrical signals spread among neighboring synapses for 10s of microns following induction of LTP at an individual synapse [55–57]. Hence, whether individual synapses are visualized in the living state before and after induction of LTP or a large number of dendrites and synapses in the path of electrical or chemical induction of LTP are evaluated, one cannot be sure exactly which synapses mediated the LTP. Others have used calcium-oxalate in an attempt to identify where synapse-specific calcium influx might have occurred following induction of LTP [58]; however, the electron dense calcium-oxalate precipitate shows high background and only those spines containing SER accumulated enough precipitate in either the control or LTP conditions. Hence, this approach to identify the potentiated synapses is biased because only about 10–15% of hippocampal dendritic spines contain SER [59]. Thus, it is not possible with current methods to define unequivocally which synapses were responsible for expressing LTP. Despite these caveats, it is reasonable to deduce that the structural changes are associated with LTP. The challenge ahead is to learn which structural changes are crucial for LTP.

6 Synaptic structure and function

Dendritic spines and postsynaptic densities

The use of ssEM has been essential to the assessment and identification of dendrites and spines that were first imaged with light microscopy either *in vivo* or in cultured neurons. GFP-labeled neurons in the rodent barrel cortex imaged *in vivo* following whisker trimming revealed an increase in the turnover of dendritic spines [12–14]. ssEM analyses of the imaged dendrites containing a GFP-directed antibody and HRP-reaction product revealed that the new spines formed synapses with presynaptic axons and that dendritic protrusions formed before synapses [12–14]. Combined light and ssEM imaging on cultured neurons revealed that Purkinje spines with synaptic contacts remain motile [60] and that induction of LTP in cultured hippocampal neurons resulted in spineogenesis that was followed by a delayed formation (>15 hours) of synapses [61]. However, the long intervals between imaging sessions could have missed synapse turnover and other studies have suggested that synapse formation on new spines could happen on a much more rapid time scale [15,62*].

Initial ssEM analyses found little or no change in synapse size or number during LTP [7,8,10]; however, warmer slice preparation and more robust statistical analyses later revealed marked and balanced change in synapse size and number during LTP in mature hippocampal slices [4**]. An increase in the turnover of spines was observed within the first 30 min after induction of LTP, as evidenced by an increase in excitatory shaft synapses, stubby spines and nonsynaptic protrusions. At 2 hours after LTP induction, small spines were lost and the remaining PSDs were enlarged (Figure 4a–e). The increase in PSD area was sufficient to compensate for the loss of spines such that the total summed PSD area per micron length of dendrite remained constant between control and LTP dendrites (Figure 4e,f). This structural synaptic scaling would have been missed with single section analyses or light microscopy, highlighting the essential need for ssEM to deduce structural plasticity, particularly in the mature hippocampus.

The ‘spine splitting’ hypothesis is another example of the requirement for ssEM to elucidate mechanisms of structural plasticity. Ultrastructural analyses found an increase in multiple synaptic boutons (MSBs) following induction of LTP in area CA1 of organotypic slices, and led to the speculation that new spines may preferentially synapse with boutons that were strongly stimulated [54]. It was also suggested that the new spines were created through a process of ‘splitting’ wherein a pre-existing spine split into two separate entities to increase synaptic density. However, ssEM and three-dimensional reconstructions revealed that the neuropil between neighboring spines from a single dendrite synapsing with the same presynaptic bouton contained numerous long axons that could not accommodate the extensive retractions and rearrangements that

would have had to occur to allow the so-called spine splitting [5]. Instead, ssEM analyses revealed that these rare same-dendrite, multi-synaptic boutons likely form via new spine outgrowth which occurs within the first 5–30 min after the onset of LTP [4**,5].


Presynaptic axons

More recently, ssEM has begun to elucidate structural changes among presynaptic axons and boutons during LTP and LTD. ssEM of organotypic slice cultures confirmed the formation of new presynaptic boutons with LTP [52] and that the varicosities that increased their turnover with LTD were indeed presynaptic boutons [51]. MSBs have been considered a potential substrate for co-activation of multiple spines and have been proposed as a mechanism to coordinate plasticity between neighboring neurons [63]. However, they occur at a relatively low frequency in stratum radiatum of area CA1 [64,65] and immuno-labeling revealed that low levels of postsynaptic glutamate receptors on spines synapsing with MSBs suggests that simultaneous depolarization would not occur [66]. Thus, while light microscopy has revealed the temporal dynamics of some presynaptic changes, ssEM was needed to correctly identify presynaptic structures.

Presynaptic boutons have large pools of vesicles that are organized into functional categories depending on their probability of release [67]. ssEM has provided an essential component by describing the distribution of vesicles within boutons of Schaffer collaterals, mossy fibers, and the neuromuscular junction [64,68,69]. Work combining FM-labeling of recycling vesicle with ssEM has been used to describe the probability of release at synapses and to reveal the distribution of recycled vesicles within boutons [68,70–72]. The combined light-level and ssEM imaging of labeled vesicles also revealed the existence of a ‘superpool’ of vesicles that are shared between neighboring boutons along an axon which expands the presynaptic release capacity well beyond the scope of individual synapses [55,73**].

Questions remain as to the fate of the ‘superpool’ vesicles with plasticity and whether the increase in presynaptic release observed with FM1-43 [39,40] corresponds to a decrease in synaptic vesicles. Single section analyses revealed a loss of vesicles within the first hour after LTP induction in both area CA1 [74] and the dentate gyrus [32], but a comprehensive three-dimensional analysis of synaptic vesicle distribution during LTP is still missing. In the lateral amygdala, ssEM did not reveal a change in the overall number of vesicles with fear conditioning, although an increase in the ratio of PSD area to docked vesicle number was observed [16]. Thus, future studies with ssEM will be necessary to examine the timing and coordination of presynaptic and postsynaptic structural changes.

Figure 4



Current Opinion in Neurobiology

Dramatic structural synaptic scaling revealed by ssEM following induction of LTP by TBS in the mature hippocampus. **(a)** EM and **(ai)** reconstruction of a small thin spine (category T, head diameter <0.45 μm). **(b)** EM and **(bi)** reconstruction of a medium thin spine (category T, head diameter >0.45 μm but <0.6 μm). **(c)** EM and **(ci)** reconstruction of a mushroom spine (M, head diameter >0.6 μm). **(d)** Small thin spines were significantly reduced in

8 Synaptic structure and function

In addition to small synaptic vesicles, ssEM has characterized the frequency and distribution of small dense core vesicles (DCVs) [75], which are known to contain active zone proteins [76]. Induction of recuperative synaptogenesis during the rewarming of acute hippocampal slices prepared with ice-cold media resulted in a loss of DCVs, suggesting that they were used to create new active zones [75].

Organelles

Local protein synthesis is required for maintenance of LTP [77] and is evidenced ultrastructurally by polyribosomes (Figure 3b,c). Polyribosomes occur every 1–2 μm along the lengths of both immature and mature hippocampal dendrites and thus intersperse among 4–10 dendritic spines [3,4^{**},6]. Single section analyses suggested a decrease in polyribosomes at the base of spines in the mature dentate gyrus at various time points after LTP [78]. In contrast, ssEM demonstrated a redistribution of polyribosomes from dendritic shafts into spines with larger PSDs at 2 hours after induction of LTP by tetanic stimulation in hippocampal slices from young [6] and mature [3] rats. More recent work found a depletion of polyribosomes that was coupled with an enlargement of PSDs at 2 hours after induction of LTP by TBS, suggesting that mRNAs may have been ‘used up’ to increase the size of the PSD [4^{**}]. Thus, ssEM is needed to detect the timing of formation and specific localization of polyribosomes.

SER enters a small percentage of spines in both the hippocampus and the lateral amygdala [17^{**},59,79]. The spine apparatus, a specialization of SER found preferentially in large spines, is involved in the regulation of calcium and the synthesis of transmembrane proteins and is important for synaptic plasticity [80]. In the lateral amygdala, ssEM revealed that fear conditioning resulted in an enlargement of PSDs on a subset of spines lacking a spine apparatus [17^{**}]. Spines that contained a spine apparatus did not further enlarge with fear conditioning, suggesting a ceiling effect, but did decrease in size with safety learning. Thus, the presence of a spine apparatus appears to have an important role in regulating changes in synaptic size in the lateral amygdala.

Recycling or sorting endosomes occur in dendritic shafts at a rate of about one per 10–20 dendritic spines [9,79]. Endosomes are involved in recycling receptors and contributing membrane to spines for enlargement with LTP. Live imaging of cultured neurons found that induction of LTP resulted in the trafficking of recycling endosomes and vesicles into spines [9]. Quantification through ssEM

of membrane surface area in these compartments within dendrites showed sufficient membrane was available to account for dendritic spine enlargement [9]. Together these findings suggest that investigation of dendritic segments through ssEM is necessary to elucidate the distribution and frequency of core dendritic resources that random sampling of individual dendritic spines would miss.

Perisynaptic astroglial processes

Characterization of perisynaptic astroglia is particularly dependent of the use of ssEM because the smallest processes are not resolved at the level of light microscopy. ssEM revealed that fewer synapses had perisynaptic astroglia in recovered slices following recuperative synaptogenesis, and that synapses were larger when perisynaptic astroglia was present [11]. Interestingly, the amount of the synaptic perimeter surrounded by perisynaptic astroglia and the distance between neighboring synapses were not proportional to synapse size, suggesting that both large and small synapses retain access to extracellular glutamate and secreted astroglial factors [11]. These findings suggest that as synapses enlarge and release more neurotransmitter, they attract astroglial processes to a discrete portion of their perimeters, further enhancing synaptic efficacy without limiting the potential for sharing extracellular resources with neighboring synapses. In addition, ssEM revealed a disruption in the three-dimensional relationship between perisynaptic astroglia and synapses in the human hippocampus removed to alleviate severe epilepsy [27].

In addition to astroglia, the interaction of microglia with synapses in the visual cortex has also been described with ssEM [81^{*}]. During normal visual experience, most microglial processes interacted with multiple synapses and preferentially localized to small and transient dendritic spines. Light deprivation decreased the motility of microglial processes and changed their localization to the vicinity of a subset of larger dendritic spines that subsequently decreased in size. These changes were reversed upon light re-exposure. Thus, in addition to the role of microglia in the immune response, they also involved in regulating spine size during visual experience.

Conclusions

In the past 60 years, ssEM has evolved into a powerful tool with which to study the ultrastructural plasticity of synapses. Advances in technology have aided the rate at which data can be acquired and analyzed, leading to an expanding interest in mapping the neurocircuitry of the

number ($P < 0.05$) while (e) PSDs on all remaining spines were significantly enlarged (** $P < 0.01$; * $P < 0.05$). (f) Reconstructions of control and LTP dendrites demonstrate equal summed PSD area per micron length of dendrite, despite large differences in their average synapse size and density. Scale cube = 0.5 μm on a side. (g) Average summed PSD area per micron length of dendrite is the same for dendrites across all times in the control and TBS-LTP conditions. Adapted from Bourne and Harris [4^{**}].

brain at the nanometer level, the so-called ‘connectome’ [82]. However, even knowing the location of every synapse in the connectome will not provide the complete answer because we need to know the functional state of each synapse, which can be further influenced by the ion channel composition in the nonsynaptic membrane of the postsynaptic dendrite or spine [83]. Detailed analysis through ssEM reveals invaluable information about the cell biological mechanisms that underlie the direction and degree of plasticity expressed by individual synapses. The finding of structural synaptic scaling during LTP emphasizes an important question of whether a few large synapses are actually more effective than many small synapses [4^{**},84]. Elucidating ultrastructural relationships between neighboring synapses will help to reveal how the weight of synaptic responses is distributed and how it is modified during plasticity and learning.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Palay SL, Palade GE: **The fine structure of neurons.** *J Biophys Biochem Cytol* 1955, **1**:69-88.
2. Gray EG: **Axo-somatic and axo-dendritic synapses of the cerebral cortex: an electron microscope study.** *J Anat* 1959, **93**:420-433.
3. Bourne JN, Sorra KE, Hurlburt J, Harris KM: **Polyribosomes are increased in spines of CA1 dendrites 2 h after the induction of LTP in mature rat hippocampal slices.** *Hippocampus* 2007, **17**:1-4.
4. Bourne JN, Harris KM: **Coordination of size and number of •• excitatory and inhibitory synapses results in a balanced structural plasticity along mature hippocampal CA1 dendrites during LTP.** *Hippocampus* 2011, **21**:354-373.

Three-dimensional analyses of dendritic segments revealed significant structural plasticity of both excitatory and inhibitory synapses following the induction of LTP with TBS. Turnover of spines within the three-dimensional analyses of dendritic segments revealed significant structural plasticity of both excitatory and inhibitory synapses following the induction of LTP with TBS. Turnover of spines within the first 30 min led to a decrease in small thin spines by 2 hours that was perfectly counterbalanced by an increase in PSD area, such that the summed PSD area per unit micron length of dendrite remained constant between control and LTP conditions. These data suggest that structural synaptic scaling may be an important cellular mechanism of plasticity in the mature hippocampus. The first 30 min led to a decrease in small thin spines by 2 hours that was perfectly counterbalanced by an increase in PSD area, such that the summed PSD area per unit micron length of dendrite remained constant between control and LTP conditions. These data suggest that structural synaptic scaling may be an important cellular mechanism of plasticity in the mature hippocampus.

5. Fiala JC, Allwardt B, Harris KM: **Dendritic spines do not split during hippocampal LTP or maturation.** *Nat Neurosci* 2002, **5**:297-298.
6. Ostroff LE, Fiala JC, Allwardt B, Harris KM: **Polyribosomes redistribute from dendritic shafts into spines with enlarged synapses during LTP in developing rat hippocampal slices.** *Neuron* 2002, **35**:535-545.
7. Popov VI, Davies HA, Rogachevsky VV, Patrushev IV, Errington ML, Gabbott PL, Bliss TV, Stewart MG: **Remodelling of synaptic morphology but unchanged synaptic density during late phase long-term potentiation (LTP): a serial section electron micrograph study in the dentate gyrus in the anaesthetised rat.** *Neuroscience* 2004, **128**:251-262.
8. Stewart MG, Medvedev NI, Popov VI, Schoepfer R, Davies HA, Murphy K, Dallerac GM, Kraev IV, Rodriguez JJ: **Chemically induced long-term potentiation increases the number of perforated and complex postsynaptic densities but does not alter dendritic spine volume in CA1 of adult mouse hippocampal slices.** *Eur J Neurosci* 2005, **21**:3368-3378.
9. Park M, Salgado JM, Ostroff L, Helton TD, Robinson CG, Harris KM, Ehlers MD: **Plasticity-induced growth of dendritic spines by exocytic trafficking from recycling endosomes.** *Neuron* 2006, **52**:817-830.
10. Sorra KE, Harris KM: **Stability in synapse number and size at 2 hr after long-term potentiation in hippocampal area CA1.** *J Neurosci* 1998, **18**:658-671.
11. Witcher MR, Kirov SA, Harris KM: **Plasticity of perisynaptic astroglia during synaptogenesis in the mature rat hippocampus.** *Glia* 2007, **55**:13-23.
12. Holtmaat A, Wilbrecht L, Knott GW, Welker E, Svoboda K: **Experience-dependent and cell-type-specific spine growth in the neocortex.** *Nature* 2006, **441**:979-983.
13. Knott GW, Holtmaat A, Wilbrecht L, Welker E, Svoboda K: **Spine growth precedes synapse formation in the adult neocortex in vivo.** *Nat Neurosci* 2006.
14. Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K: **Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex.** *Nature* 2002, **420**:788-794.
15. Zito K, Knott G, Shepherd GM, Shenolikar S, Svoboda K: **Induction of spine growth and synapse formation by regulation of the spine actin cytoskeleton.** *Neuron* 2004, **44**:321-334.
16. Ostroff LE, Cain CK, Jindal N, Dar N, Ledoux JE: **Stability of presynaptic vesicle pools and changes in synapse morphology in the amygdala following fear learning in adult rats.** *J Comp Neurol* 2011.
17. Ostroff LE, Cain CK, Bedont J, Monfils MH, Ledoux JE: **Fear and •• safety learning differentially affect synapse size and dendritic translation in the lateral amygdala.** *Proc Natl Acad Sci U S A* 2010, **107**:9418-9423.

Following fear or safety learning, dendrites from the lateral amygdala were analyzed using ssEM. PSDs on spines without a spine apparatus increased in size with fear learning while spines with a spine apparatus decreased in size with safety learning. In addition, polyribosomes and multivesicular bodies were increased throughout dendrites after fear conditioning. These data demonstrate that structural synaptic plasticity and redistribution of dendritic resources are core features of fear and safety learning in the lateral amygdala.

18. Sosinsky GE, Crum J, Jones YZ, Lanman J, Smarr B, Terada M, • Martone ME, Deerinck TJ, Johnson JE, Ellisman MH: **The combination of chemical fixation procedures with high pressure freezing and freeze substitution preserves highly labile tissue ultrastructure for electron tomography applications.** *J Struct Biol* 2008, **161**:359-371.

This study found that combining chemical fixation with high pressure freezing resulted in superior tissue preservation than chemical fixation alone. In particular, fragile tissue such as the brain shows excellent preservation with the combined techniques. These data demonstrate a continuing evolution of methods for enhancing the use of electron microscopy in structural synaptic plasticity.

19. Rostaing P, Real E, Siksou L, Lechaire JP, Boudier T, Boeckers TM, Gertler F, Gundelfinger ED, Triller A, Marty S: **Analysis of synaptic ultrastructure without fixative using high-pressure freezing and tomography.** *Eur J Neurosci* 2006, **24**:3463-3474.
20. Reymann KG, Frey JU: **The late maintenance of hippocampal LTP: requirements, phases, ‘synaptic tagging’, ‘late-associativity’ and implications.** *Neuropharmacology* 2007, **52**:24-40.
21. Burgoyne RD, Gray EG, Sullivan K, Barron J: **Depolymerization of dendritic microtubules following incubation of cortical slices.** *Neurosci Lett* 1982, **31**:81-85.
22. Feig S, Lipton P: **N-methyl-D-aspartate receptor activation and Ca²⁺ account for poor pyramidal cell structure in hippocampal slices.** *J Neurochem* 1990, **55**:473-483.

10 Synaptic structure and function

23. Fiala JC, Kirov SA, Feinberg MD, Petrak LJ, George P, Goddard CA, Harris KM: **Timing of neuronal and glial ultrastructure disruption during brain slice preparation and recovery in vitro.** *J Comp Neurol* 2003, **465**:90-103.
24. Sajikumar S, Navakkode S, Frey JU: **Protein synthesis-dependent long-term functional plasticity: methods and techniques.** *Curr Opin Neurobiol* 2005.
25. Ullian EM, Sapperstein SK, Christopherson KS, Barres BA: **Control of synapse number by glia.** *Science* 2001, **291**:657-661.
26. Ullian EM, Christopherson KS, Barres BA: **Role for glia in synaptogenesis.** *Glia* 2004, **47**:209-216.
27. Witcher MR, Park YD, Lee MR, Sharma S, Harris KM, Kirov SA: **Three-dimensional relationships between perisynaptic astroglia and human hippocampal synapses.** *Glia* 2010, **58**:572-587.
28. Kirov SA, Sorra KE, Harris KM: **Slices have more synapses than perfusion-fixed hippocampus from both young and mature rats.** *J Neurosci* 1999, **19**:2876-2886.
29. Kirov SA, Petrak LJ, Fiala JC, Harris KM: **Dendritic spines disappear with chilling but proliferate excessively upon rewarming of mature hippocampus.** *Neuroscience* 2004, **127**:69-80.
30. Roelandse M, Matus A: **Hypothermia-associated loss of dendritic spines.** *J Neurosci* 2004, **24**:7843-7847.
31. Bourne JN, Kirov SA, Sorra KE, Harris KM: **Warmer preparation of hippocampal slices prevents synapse proliferation that might obscure LTP-related structural plasticity.** *Neuropharmacology* 2007, **52**:55-59.
32. Fifkova E, Van Harrevelde A: **Long-lasting morphological changes in dendritic spines of dentate granular cells following stimulation of the entorhinal area.** *J Neurocytol* 1977, **6**:211-230.
33. Van Harrevelde A, Fifkova E: **Swelling of dendritic spines in the fascia dentata after stimulation of the perforant fibers as a mechanism of post-tetanic potentiation.** *Exp Neurol* 1975, **49**:736-749.
34. Trommald M, Vaaland JL, Blackstad TW, Andersen P: **Dendritic spine changes in rat dentate granule cells associated with long-term potentiation.** In *Neurotoxicity of Excitatory Amino Acids*. Edited by Guidotti A, Costa E. New York: Raven Press, Ltd.; 1990:163-174.
35. Trommald M, Hulleberg G, Andersen P: **Long-term potentiation is associated with new excitatory spine synapses on rat dentate granule cells.** *Learn Mem* 1996, **3**:218-228.
36. Lee KS, Schottler F, Oliver M, Lynch G: **Brief bursts of high-frequency stimulation produce two types of structural change in rat hippocampus.** *J Neurophysiol* 1980, **44**:247-258.
37. Chang FL, Greenough WT: **Transient and enduring morphological correlates of synaptic activity and efficacy change in the rat hippocampal slice.** *Brain Res* 1984, **309**:35-46.
38. Kang H, Welcher AA, Shelton D, Schuman EM: **Neurotrophins and time: different roles for TrkB signaling in hippocampal long-term potentiation.** *Neuron* 1997, **19**:653-664.
39. Bayazitov IT, Richardson RJ, Fricke RG, Zakharenko SS: **Slow presynaptic and fast postsynaptic components of compound long-term potentiation.** *J Neurosci* 2007, **27**:11510-11521.
40. Zakharenko SS, Zablou L, Siegelbaum SA: **Visualization of changes in presynaptic function during long-term synaptic plasticity.** *Nat Neurosci* 2001, **4**:711-717.
41. Morgan SL, Teyler TJ: **Electrical stimuli patterned after the theta-rhythm induce multiple forms of LTP.** *J Neurophysiol* 2001, **86**:1289-1296.
42. White JG, Southgate E, Thomson JN, Brenner S: **Factors that determine connectivity in the nervous system of *Caenorhabditis elegans*.** *Cold Spring Harb Symp Quant Biol* 1983, **48**(Pt 2):633-640.
43. Mishchenko Y, Hu T, Spacek J, Mendenhall J, Harris KM, ●● Chklovskii DB: **Ultrastructural analysis of hippocampal neuropil from the connectomics perspective.** *Neuron* 2010, **67**:1009-1020.
- Automated dense reconstructions of serial sections from rat hippocampus revealed that axo-dendritic touches often failed to predict a synapse. The largest touch between an axonal bouton and spine identified sites of synaptic contact with relatively high accuracy, but still missed a large fraction of synapses. This paper highlights the potential pitfalls of a connectomic approach that disregards ultrastructural details.
44. Bock DD, Lee WC, Kerlin AM, Andermann ML, Hood G, ●● Wetzel AW, Yurgenson S, Soucy ER, Kim HS, Reid RC: **Network anatomy and in vivo physiology of visual cortical neurons.** *Nature* 2011, **471**:177-182.
- Two-photon calcium imaging was combined with ssEM to elucidate the relationship between structure and function of neurons in mouse primary visual cortex. The data revealed that inhibitory interneurons received convergent anatomical input from excitatory neurons that exhibited a broad range of preferred orientations. This paper demonstrates a direct link between the functional output of neurons and their synaptic connectivity at the ultrastructural level.
45. Helmstaedter M, Briggman KL, Denk W: **High-accuracy neurite reconstruction for high-throughput neuroanatomy.** *Nat Neurosci* 2011, **14**:1081-1088.
46. Geinisman Y, de Toledo-Morrell L, Morrell F, Heller RE, Rossi M, Parshall RF: **Structural synaptic correlate of long-term potentiation: formation of axospinous synapses with multiple, completely partitioned transmission zones.** *Hippocampus* 1993, **3**:435-445.
47. Desmond NL, Levy WB: **Changes in the postsynaptic density with long-term potentiation in the dentate gyrus.** *J Comp Neurol* 1986, **253**:476-482.
48. O'Malley A, O'Connell C, Regan CM: **Ultrastructural analysis reveals avoidance conditioning to induce a transient increase in hippocampal dentate spine density in the 6 hour post-training period of consolidation.** *Neuroscience* 1998, **87**:607-613.
49. Eyre MD, Richter-Levin G, Avital A, Stewart MG: **Morphological changes in hippocampal dentate gyrus synapses following spatial learning in rats are transient.** *Eur J Neurosci* 2003, **17**:1973-1980.
50. Rusakov DA, Davies HA, Harrison E, Diana G, Richter-Levin G, Bliss TV, Stewart MG: **Ultrastructural synaptic correlates of spatial learning in rat hippocampus.** *Neuroscience* 1997, **80**:69-77.
51. Becker N, Wierenga CJ, Fonseca R, Bonhoeffer T, Nagerl UV: **LTD induction causes morphological changes of presynaptic boutons and reduces their contacts with spines.** *Neuron* 2008, **60**:590-597.
52. Nikonenko I, Jourdain P, Muller D: **Presynaptic remodeling contributes to activity-dependent synaptogenesis.** *J Neurosci* 2003, **23**:8498-8505.
53. Toni N, Buchs PA, Nikonenko I, Povilaitite P, Parisi L, Muller D: **Remodeling of synaptic membranes after induction of long-term potentiation.** *J Neurosci* 2001, **21**:6245-6251.
54. Toni N, Buchs PA, Nikonenko I, Bron CR, Muller D: **LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite.** *Nature* 1999, **402**:421-425.
55. Darcy KJ, Staras K, Collinson LM, Goda Y: **Constitutive sharing of recycling synaptic vesicles between presynaptic boutons.** *Nat Neurosci* 2006, **9**:315-321.
56. Harvey CD, Yasuda R, Zhong H, Svoboda K: **The spread of Ras activity triggered by activation of a single dendritic spine.** *Science* 2008, **321**:136-140.
57. Rose J, Jin SX, Craig AM: **Heterosynaptic molecular dynamics: locally induced propagating synaptic accumulation of CaM kinase II.** *Neuron* 2009, **61**:351-358.
58. Buchs PA, Muller D: **Induction of long-term potentiation is associated with major ultrastructural changes of activated synapses.** *Proc Natl Acad Sci U S A* 1996, **93**:8040-8045.
59. Spacek J, Harris KM: **Three-dimensional organization of smooth endoplasmic reticulum in hippocampal CA1 dendrites and dendritic spines of the immature and mature rat.** *J Neurosci* 1997, **17**:190-203.

60. Dunaevsky A, Blazeski R, Yuste R, Mason C: **Spine motility with synaptic contact.** *Nat Neurosci* 2001, **4**:685-686.
61. Nagerl UV, Kostinger G, Anderson JC, Martin KA, Bonhoeffer T: **Protracted synaptogenesis after activity-dependent spinogenesis in hippocampal neurons.** *J Neurosci* 2007, **27**:8149-8156.
62. Zito K, Scheuss V, Knott G, Hill T, Svoboda K: **Rapid functional maturation of nascent dendritic spines.** *Neuron* 2009, **61**:247-258. Hippocampal spines were analyzed using two-photon glutamate uncaging, whole-cell recording, Ca^{2+} imaging, and EM to determine the time course of spine morphogenesis and synapse formation. New spines (<2.5 hours) were found to express AMPA and NMDA-mediated currents that were comparable to mature spines, suggesting that new spines can rapidly form functional synapses.
63. Harris KM: **How multiple-synapse boutons could preserve input specificity during an interneuronal spread of LTP.** *Trends Neurosci* 1995, **18**:365-369.
64. Shepherd GM, Harris KM: **Three-dimensional structure and composition of CA3 — CA1 axons in rat hippocampal slices: implications for presynaptic connectivity and compartmentalization.** *J Neurosci* 1998, **18**:8300-8310.
65. Sorra KE, Harris KM: **Occurrence and three-dimensional structure of multiple synapses between individual radiatum axons and their target pyramidal cells in hippocampal area CA1.** *J Neurosci* 1993, **13**:3736-3748.
66. Nicholson DA, Geinisman Y: **Axospinous synaptic subtype-specific differences in structure, size, ionotropic receptor expression, and connectivity in apical dendritic regions of rat hippocampal CA1 pyramidal neurons.** *J Comp Neurol* 2009, **512**:399-418.
67. Denker A, Rizzoli SO: **Synaptic vesicle pools: an update.** *Front Synaptic Neurosci* 2010, **2**:135.
68. Richards DA, Guatimosim C, Rizzoli SO, Betz WJ: **Synaptic vesicle pools at the frog neuromuscular junction.** *Neuron* 2003, **39**:529-541.
69. Chicurel ME, Harris KM: **Serial electron microscopy of CA3 dendritic spines synapsing with mossy fibers of rat hippocampus.** *Soc Neurosci Abs* 1989, **15**:256.
70. Schikorski T, Stevens CF: **Morphological correlates of functionally defined synaptic vesicle populations.** *Nat Neurosci* 2001, **4**:391-395.
71. Branco T, Marra V, Staras K: **Examining size-strength relationships at hippocampal synapses using an ultrastructural measurement of synaptic release probability.** *J Struct Biol* 2010, **172**:203-210.
72. Rizzoli SO, Betz WJ: **The structural organization of the readily releasable pool of synaptic vesicles.** *Science* 2004, **303**:2037-2039.
73. Staras K, Branco T, Burden JJ, Pozo K, Darcy K, Marra V, Ratnayaka A, Goda Y: **A vesicle superpool spans multiple presynaptic terminals in hippocampal neurons.** *Neuron* 2010, **66**:37-44.
- Fluorescence imaging, correlative EM, and modeling was used to demonstrate that some recycling pool vesicles form part of a larger 'superpool' that moves between multiple presynaptic boutons. The 'superpool' vesicles are highly mobile and their exchange at synapses is regulated in part by a TrkB-receptor-dependent mechanism. These data suggest a potentially important role in modifying the composition of synaptic vesicle pools during plasticity.
74. Applegate MD, Kerr DS, Landfield PW: **Redistribution of synaptic vesicles during long-term potentiation in the hippocampus.** *Brain Res* 1987, **401**:401-406.
75. Sorra KE, Mishra A, Kirov SA, Harris KM: **Dense core vesicles resemble active-zone transport vesicles and are diminished following synaptogenesis in mature hippocampal slices.** *Neuroscience* 2006, **141**:2097-2106.
76. Tao-Cheng JH: **Activity-related redistribution of presynaptic proteins at the active zone.** *Neuroscience* 2006, **141**:1217-1224.
77. Steward O, Schuman EM: **Protein synthesis at synaptic sites on dendrites.** *Annu Rev Neurosci* 2001, **24**:299-325.
78. Desmond NL, Levy WB: **Morphological correlates of long-term potentiation imply the modification of existing synapses, not synaptogenesis, in the hippocampal dentate gyrus.** *Synapse* 1990, **5**:139-143.
79. Cooney JR, Hurlburt JL, Selig DK, Harris KM, Fiala JC: **Endosomal compartments serve multiple hippocampal dendritic spines from a widespread rather than a local store of recycling membrane.** *J Neurosci* 2002, **22**:2215-2224.
80. Deller T, Orth CB, Del TD, Vlachos A, Burbach GJ, Drakew A, Chabanis S, Korte M, Schwegler H, Haas CA, Frotscher M: **A role for synaptopodin and the spine apparatus in hippocampal synaptic plasticity.** *Ann Anat* 2007, **189**:5-16.
81. Tremblay ME, Lowery RL, Majewska AK: **Microglial interactions with synapses are modulated by visual experience.** *PLoS Biol* 2010, **8**:e1000527. Microglia in the visual cortex were analyzed using two-photon *in vivo* imaging, immuno-EM and ssEM with three-dimensional reconstructions. Microglial processes were found to be dynamic and associate with small or transient spines during visual experience and became less motile and associated with larger spines that subsequently decreased in size during light deprivation. These data suggest a role for microglia in the modification of synaptic circuitry even in the healthy brain.
82. Lichtman JW, Sanes JR: **Ome sweet ome: what can the genome tell us about the connectome?** *Curr Opin Neurobiol* 2008, **18**:346-353.
83. Kim J, Jung SC, Clemens AM, Petralia RS, Hoffman DA: **Regulation of dendritic excitability by activity-dependent trafficking of the A-type K^+ channel subunit Kv4.2 in hippocampal neurons.** *Neuron* 2007, **54**:933-947.
84. Bourne J, Harris KM: **Do thin spines learn to be mushroom spines that remember?** *Curr Opin Neurobiol* 2007.